

DECIPHERING THE COSMIC STAR FORMATION HISTORY AND THE NATURE OF TYPE IA SUPERNOVAE BY FUTURE SUPERNOVA SURVEYS

TAKESHI ODA AND TOMONORI TOTANI

Department of Astronomy, School of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, JAPAN

Draft version February 2, 2008

ABSTRACT

We investigate the prospects of future supernova searches to get meaningful constraints about the cosmic star formation history (CSFH) and the delay time of type Ia supernovae from star formation (τ_{Ia}), based only on supernova data. Here we parameterize the CSFH by two parameters, α and β that are the evolutionary indices [$\propto (1+z)^{\{\alpha, \beta\}}$] at $z \gtrsim 1$ and $\lesssim 1$, respectively, and quantitatively examined how well the three parameters (α , β , and τ_{Ia}) can be constrained in ongoing and future supernova surveys. We found that the type classification of detected supernovae down to the magnitude of $I_{AB} \sim 27$ is essential, to get useful constraint on β . The parameter τ_{Ia} can also be constrained within an accuracy of $\sim 1\text{--}2$ Gyr, without knowing α that is somewhat degenerate with τ_{Ia} . This might be potentially achieved by ground-based surveys but depending on the still highly uncertain type-classification by imaging data. More reliable classification will be achieved by the *SNAP* mission. The supernova counts at a magnitude level of I_{AB} or $K_{AB} \sim 30$ will allow us to break degeneracies between α and τ_{Ia} and independently constrain all the three parameters, even without knowing supernova types. This can be achieved by the *SNAP* and *JWST* missions, having different strength of larger statistics and reach to higher redshifts, respectively. The dependence of observable quantities on survey time intervals is also quantitatively calculated and discussed.

Subject headings: cosmology:observations — supernova:general

1. INTRODUCTION

It is important to measure the cosmic supernova (SN) rate at present as well as at high redshift, to understand the cosmic star formation history (CSFH) and to get a hint for the origin of the type Ia supernovae (SNe Ia). Although the CSFH up to $z \sim 5$ has been intensively studied based on star formation rate (SFR) indicators of high- z galaxies (Lilly et al. 1996; Madau et al. 1996; see Totani & Takeuchi 2002 for a recent compilation of observed data by various methods), the derived CSFH is still a matter of debate since the estimated SFR densities are considerably different depending on authors and methods. The CSFH inferred from SN rate evolution includes completely independent and complementary information. CSFH estimated by galactic SFR, especially at higher redshifts, is inevitably biased toward bright galaxies; in many studies SFR was corrected by extrapolating the galaxy luminosity function to the highly uncertain faint end, well below the observational limit. However, SNe in any galaxy including dwarfs, or even those in truly intergalactic space, can equally be detected by SN surveys and hence it is completely free from this bias. The galactic SFR indicators and core-collapse SN rate generally trace the massive star formation, but the SN Ia rate gives some information for low or intermediate mass (typically a few solar mass) star formation (e.g., Yungelson & Livio 2000). The delay time of SNe Ia from star formation to the explosion may be inferred from observations, giving a useful hint for the still unknown progenitors of SNe Ia.

There are a number of publications about theoretical predictions of high- z SN rate as a probe of CSFH and the progenitor of SNe Ia (Totani et al. 1996; Madau et al.

1998; Yungelson & Livio 1998, 2000; Gilliland et al. 1999; Sullivan et al. 2000b; Kobayashi et al. 2000). In these papers, only the time evolution of the cosmic SN rate density, which is simply converted from CSFH, was shown. However, to compare with observed data it is necessary to predict the expected number of SNe limited by a given sensitivity of an observation. A work in this direction was first presented by Dahlén & Fransson (1999, hereafter DF99).

On the other hand, in recent years there was a remarkable progress about measurement of type Ia and core-collapse supernova rate, pushing out the maximum distance from $z \sim 0$ to ~ 1.0 (Pain et al. 1996; Cappellaro et al. 1999; Hardin et al. 2000; Gal-Yam et al. 2002; Madgwick et al. 2003; Gal-Yam & Maoz 2004; Dahlén et al. 2004). Some interesting implications on the CSFH and the Ia delay time have already been obtained by analyses of these data (Pain et al. 2002; Gal-Yam et al. 2003; Maoz & Gal-Yam 2004; Blanc et al. 2004; Strolger et al. 2004), though statistics is not large enough to derive decisive conclusions. For example, one must use information on CSFH derived by other methods than SNe to derive a strong constraint on the Ia delay time. Future SN surveys will provide much better constraints and may allow to determine the interesting parameters only by supernova information.

However, the cosmic SN rate evolution is dependent on a number of parameters that are often degenerate with each other, and it is still unclear how to get useful information from future SN surveys. The purpose of this paper is to examine what can be learn from future SN surveys, and find the best strategy of analyzing data to get useful information. We set our goal as to constrain the three interesting parameters only based on informa-

tion obtained by supernova surveys: the rate of increase of the cosmic SFR from $z = 0$ to ~ 1 , the evolution of the SFR beyond $z \sim 1$, and the SN Ia delay time, by breaking degeneracy between these parameters. We will make predictions on various observational quantities for future surveys and discuss how well these parameters can be constrained in the near future.

SNe are detected by variability in more than two epochs of observations. If a reference frame that is temporally well separated from new observations is available, we can measure the real SN flux relative to the reference frame. However, the first SN candidates are often selected from a flux variability on a shorter time baseline that is comparable to the expected SN time scale (i.e., month), to find SNe at early brightening phase, and to avoid contamination of active galactic nuclei (AGNs) having longer variability time scales. The latest deep SN survey by GOODS was also based on a short time variability. This selection process should have some effect on the event rate and hence on SN rate studies, even though the real flux and light curve can be measured by using the reference frame in a distant past. Therefore, a prediction of expected SN counts as a function of variability flux limits for a given observing time interval, rather than as a function of the real SN flux in a single-epoch snapshot, would be useful to quantitatively understand the dependence of the event rate on the variability time scale. This was not quantitatively presented in earlier publications, and here we will present predictions in such a form and discuss possible effects of choosing different observing intervals.

The plan of this paper is as follows. In §2 we describe basic formulations and calculation methods, and in §3 we present the results of SN detection rate with some discussions on the choice of the observing intervals. In §4 we propose the strategy to decipher the CSFH and the Ia delay time, and discuss the feasibility of our strategy referencing to the plans of future proposed SN surveys. Discussions and conclusions will be presented in §5. Throughout this paper, the standard Λ CDM universe is assumed with following values of the cosmological parameters: $\Omega_M = 0.3$, $\Omega_\Lambda = 0.7$, $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The AB magnitude system is used for the zero-point of photometry, unless otherwise stated.

2. FORMULATIONS AND CALCULATION METHODS

2.1. Basic Equations

There are a variety of supernova searches with different search methods and sampling times, but the selection process of supernovae can essentially be considered as a search of transient objects with a sensitivity magnitude limit to flux variability, m_{lim} , between two epochs separated by a time interval T . In reality a more complicated selection procedures (e.g., more than two epochs) may be taken, but it can be expressed by a combination of multiple two-epoch selections. The total number of detectable SNe, $N(m_{\text{lim}}, T)$, in a survey observing an area of Ω_{obs} is calculated from the following equation:

$$N(m_{\text{lim}}, T) = \sum_i \int_0^\infty \frac{dN_i(m_{\text{lim}}, T, z)}{dz} dz, \quad (1)$$

where

$$\frac{dN_i(m_{\text{lim}}, T, z)}{dz} = \Omega_{\text{obs}} \frac{r_i(z)}{1+z} \frac{dV(z)}{dz} f_i(z, m_{\text{lim}}, T). \quad (2)$$

Here the subscript i denotes supernova types and the summation in Eq. (1) is over all the types of SNe. The comoving density of SN rate is described by $r_i(z)$, where the rate is measured by the cosmic time (or the rest-frame time at z). The factor of $(1+z)^{-1}$ accounts for the cosmological time dilation. The comoving volume element $dV(z)/dz$ per unit solid angle is in the ordinary definition, and $f_i(z, m_{\text{lim}}, T)$ is a function called effective visibility time (EVT) or control time, which is defined by the observer's time (at $z = 0$). This function calculates a time duration in which a SN can be detected, and it depends on the spectra and light curves as well as the magnitude limit, observing interval, and redshift. In a prediction for a snapshot observation without taking into account variability, EVT is a duration of a part of SN light curve when a supernova is brighter than the magnitude limit. On the other hand, in a variability-limited survey with an interval of T , the EVT is the time duration when the variability flux is brighter than the magnitude limit for variability, i.e.,

$$f_i = \int H(m_{\text{lim}} - m_{\text{var}}) dt, \quad (3)$$

where t is a time coordinate for supernova evolution and $H(x)$ is the step function [$H(x) = 1$ and 0 for $x \geq 0$ and < 0 , respectively], and

$$m_{\text{var}}(z, t, T) = -2.5 \log \left\{ \left| 10^{-0.4m(z,t)} - 10^{-0.4m(z,t+T)} \right| \right\} \quad (4)$$

is the magnitude of the variability flux between two epochs of t and $t+T$. Here $m(z, t)$ is the real magnitude of a SN in a given waveband at redshift z and epoch t . It should be noted that t is the time for observer and the cosmological time dilation must be included in calculation of $m(z, t)$.

2.2. Light Curves and Spectra of SNe

The apparent SN magnitude $m(z, t)$ can be calculated if redshift, light curve, and spectral evolution are known, taking into account the K-correction and cosmological time dilation by a standard manner. We mostly followed the prescription by DF99 for the calculation of supernova light curves and spectra, but we collected and used the latest data published after 1999 if available, to make the most reliable prediction made so far. Here we only describe these new data, and other ingredients that are not mentioned here are the same with DF99. The lightcurves and blackbody temperatures used in our calculation are shown in Figure 1.

The spectra and light curves are taken from Nugent et al. (2002) for type Ia SNe, while those of SNe 1999em (Hamuy et al. 2001), 1979c (Branch et al. 1981), and 1995G (Pastorello et al. 2002) are used for types IIP, IIL, and IIn, respectively. The light curves of type Ib and Ic SNe are similar, and we treat these two as a single population, for which the light curve model is constructed based on the data of 1994I, 1999ex and 2002ap (Richmond et al. 1996; Stritzinger et al. 2002; Foley et al. 2003). The peak magnitudes differ considerably from supernova to supernova even within a

type, and this is taken into account by integrating the peak luminosity function of SNe for all types, assuming a Gaussian form in magnitude with mean magnitudes and dispersions taken from local values measured by Richardson et al. (2002) (shown in Table 1). We use the same relative proportions of the four subclasses of core-collapse SNe with DF99, which are assumed to be constant with the cosmic time. The sum of the relative proportions is not unity because we do not include peculiar SN1987A-type supernovae, whose peak magnitude is much fainter ($M_B \sim -15$) and hence they contribute little to detection number of high- z SNe. For SNe Ia, the well-known peak-luminosity and lightcurve-shape relation (Phillips 1993; Phillips et al. 1999; Perlmutter et al. 1997; Strolger et al. 2004) is taken into account, while the same shape of lightcurves are used for core-collapse SNe for all peak magnitudes. The color/lightcurve-shape relation of SNe Ia (Phillips et al. 1999) is not included in our calculation. We found that this effect would hardly change the results presented in our paper.

We do not take account of the Galactic extinction, which can be easily corrected by the known extinction map. On the other hand, extinction in host galaxies should be dependent on their various properties and location of SNe in them, and there should be a large variety (Hatano et al. 1998; Totani & Kobayashi 1999). Since it is difficult to construct a realistic model for this, we simply assume a mean reddening of $E(B - V) = 0.1$ and the standard Milky-Way type extinction curve (Cardelli et al. 1989) to make our prediction more realistic than no extinction case. (A detailed discussion on this issue will be given in Section 5.)

2.3. the Cosmic Star Formation History

The CSFH must be specified to predict SN rate as a function of redshift. Here we use a phenomenological CSFH model (Gal-Yam & Maoz 2004) inferred from recent observations of high- z galaxies:

$$\Phi(z) = \frac{2^{0.2} \Phi(1.2)}{\left[\left(\frac{2.2}{1+z} \right)^{5\alpha} + \left(\frac{2.2}{1+z} \right)^{5\beta} \right]^{0.2}}. \quad (5)$$

Here α and β are the SFR evolutionary indices at high ($z \gtrsim 1$) and low ($z \lesssim 1$) redshift, respectively. There is a consensus that the cosmic SFR increases from $z = 0$ to $\sim 1-2$, while the estimated values differ as $\beta \sim 1-3$ (e.g., Lilly et al. 1996; Sullivan et al. 2000a; Wilson et al. 2002; Lanzetta et al. 2002; Schiminovich et al. 2005). It is even more uncertain whether the SFR density is roughly constant to $z \sim 5$ or declines beyond $z \sim 1-2$ (e.g., Madau et al. 1996; Hughes et al. 1998). In our model, a cut off at high redshift $z_u = 10$ is introduced, beyond which there is no star formation, though this cut off hardly affects our results.

The supernova rate density is calculated by the usual manner as in earlier papers (Madau et al. 1998, DF99). The rate of core-collapse SNe (all types except Ia) is simply proportional to the CSFH, but type Ia rate is converted from the CSFH by convolving the delay function, i.e., the distribution function of the delay time t_D from star formation to the type Ia events¹. We mainly

use a standard exponential form for the delay function, $f_D(t_D) \propto \exp(-t_D/\tau_{\text{Ia}})$ (e.g. Madau et al. 1998; Gal-Yam & Maoz 2004; Strolger et al. 2004) in this paper. Only in §4.1 we calculate using the Gaussian type function defined Strolger et al. (2004) to check the difference between them.

In most earlier papers, the normalization of SN rate history is determined by the number of supernova events from a unit mass of star formation, but this is highly uncertain. To avoid this uncertainty, in this paper we always renormalize the calculated SN rate history to the local SN rate measured by Cappellaro et al. (1999). This prescription is also adequate for our purpose, i.e., deciphering CSFH and τ_{Ia} only based on supernova data. The renormalization is done separately for type Ia and core-collapse SNe, whose local rate density is 3.0×10^{-5} and $7.5 \times 10^{-5} \text{ yr}^{-1} \text{ Mpc}^{-3}$, respectively (Cappellaro 2003). Therefore the SN detection number calculated in this paper can be considered as a ratio to the local SN rates. It should be noted that, with the formulations above, the functional form of stellar initial mass function (IMF) does not affect the SN rate calculation, provided that the IMF is universal and constant with time.

3. THE SUPERNOVA DETECTION RATE AND OBSERVING TIME INTERVALS

In this section we present some results on the expected SN detection rate. To see the dependence on the model parameters that we are interested in, we use two CSFH models and two values of the Ia delay time, which are largely different but still in the reasonable range. For one CSFH model, called the “peak” model, $(\alpha, \beta) = (-1, 3)$ are used, while $(\alpha, \beta) = (0, 1)$ are used for the other, called the “flat” model. The Ia delay time is set to $\tau_{\text{Ia}} = 0.3$ and 3.0 Gyr, which are referred as “short” and “long” models, respectively.

The left panel of Fig.2 shows the total number of detectable SNe as a function of the magnitude limit for variability flux, which is a similar plot to popular galaxy counts or star counts as functions of magnitude limit. (Therefore we call this plot as “SN counts”, hereafter.) The SN counts depend on the interval of observation; shorter intervals predict less number of detectable SNe. The long baseline case ($\gtrsim 360$ days) correspond to the early studies like DF99, and our results are mostly consistent with those of DF99.

The right panel of figure 2 shows the ratio of SN counts with different intervals of 30 and 360 days, $N(30)/N(360)$, for different models of the SN rate history. The dependence of this ratio on the SN rate history is not large [$\Delta\{N(30)/N(360)\} \lesssim 0.03$].

When SNe are selected by flux variability in two epochs of observations, SNe can be classified into two categories: SNe whose second-epoch flux increases or decreases compared with the first epoch. We call these as brightening and fading SNe. (Therefore it is different from the derivative of light curves at a given time.) The ratio of number of brightening to fading SNe, N_B/N_F , is an observable

time from the end of the main-sequence phase to the occurrence of SNe Ia. However, such definition is dependent on stellar IMF, and here we define it from the time of star formation. For a typical IMF (the Salpeter’s function with the cut-offs at 0.1 and 100 $M_\odot$), the mean difference between the two definitions weighted over stellar mass is about 0.3 Gyr.

¹ In some earlier publications, the delay time t_D is defined as the

quantity that can easily be obtained in any supernova surveys, and which is shown in Figure 3. This ratio is determined by the observing interval and shape of SN light curves, and potentially it could give some information of SN type mix. When the two images are separated enough, both the number of the brightening and fading phase should be the same, and the ratio becomes unity. For a fixed interval, the ratio becomes smaller at deeper magnitude limit, because the faint and long fading phase of relatively close supernovae mainly contribute to N_F . On the other hand, the ratio becomes larger than the unity at the brightest limiting magnitude in short interval observations. In this case only SNe having large variability can be detected, and the initial rapid brightening phase contributes significantly. Because of the combined effects of long decaying phase and cosmological time dilation, longer intervals are required in deeper observations, to get two independent reference frames [$N_B/N_F = 1$]. One year interval is enough for this in most cases, but it starts to be insufficient for the deepest surveys possible in the near future ($I_{AB} \gtrsim 30$).

The right panel of figure 3 shows change of N_B/N_F by different CSFH models and/or type Ia delay time. The difference is not particularly large and it seems difficult to use the observational quantities of $N(30)/N(360)$ or N_B/N_F to discriminate different SN rate histories. However, these quantities change significantly with the survey depth and time intervals, which can easily be measured by observations. Therefore comparing these predictions with actual data will provide a useful consistency check, such as examination of possible contamination by non-SN events like AGNs.

4. DECIPHERING THE CSFH AND THE Ia DELAY TIME

In the framework described above, the three parameters that we want to constrain by future observations are: (1) the low-redshift CSFH index, β , (2) the high-redshift CSFH index, α , and (3) the Ia delay time, τ_{Ia} . The question that we want to answer is whether we can constrain these three parameters by future supernova data, breaking the degeneracy among them. For this purpose we try the following strategy. It is expected that, in relatively shallow surveys, only supernovae at $z \lesssim 1$ are detectable and the α parameter does not affect significantly the observable quantities, allowing us to constrain mainly β and τ_{Ia} . (The size of the effect of changing α will quantitatively be checked later.) The survey time interval is set to 30 days in all the results shown in this section, and our main conclusions do not depend much on the choice of the time interval.

4.1. Determining β and τ_{Ia} by SNe at $z \lesssim 1$

Then our first approach is to constrain β and τ_{Ia} by two independent observational quantities at relatively shallow supernova surveys. The quantity that is the most easily obtained is the total detected number of SNe including all types at the detection limit of a survey. For the second observational quantity, we examine the redshift distribution and the fraction of type Ia supernovae (f_{Ia}). Redshift distribution is easier to obtain by photometric redshifts of host galaxies without spectroscopy, than f_{Ia} that requires spectroscopy for a firm determination. However, we find that there is a considerable degeneracy between β and τ_{Ia} in the redshift distribu-

tion, and the type Ia fraction has a stronger power to constrain β and τ_{Ia} . We use $\alpha = 0$ for the results presented in this section.

4.1.1. Redshift Distribution

In order to see the power of breaking degeneracy by redshift distribution, we calculate the redshift distributions and type Ia fractions using two cases where the values of β and τ_{Ia} are very different but the total expected number of SN detection is the same. Figure 4 shows the results, where the used parameters are $(\beta, \tau_{Ia}[\text{Gyr}]) = (3.0, 4.0)$ and $(2.3, 0.3)$. It can be seen that there is only a small difference in the redshift distribution of SNe of all types. The mean redshift differs only by ~ 0.2 , and it is comparable with the accuracy of photometric redshifts of galaxies (e.g. Furusawa et al. 2000). Even if accurate spectroscopic redshifts are available, such a small difference of distribution may be hidden by uncertainties in theoretical modeling.

Gal-Yam & Maoz (2004, hereafter GM04) investigated how the three parameters (α, β, τ_{Ia}) can be constrained by the redshift distribution of about 1,000 SNe in future surveys, and they concluded that the redshift distribution gives some useful constraint, which appears somewhat different from our findings here. In fact, there is no discrepancy; we made a similar analysis to GM04, and confirmed that our results are mostly similar to those in GM04. Here we note that GM04 checked only one parameter set, $(\alpha, \beta, \tau_{Ia}) = (-2, 4, 1)$, as the input parameters for simulations. On the other hand, we found the above degeneracy of the redshift distribution by using smaller values of β . This suggests that the degeneracy becomes more serious than found by GM04 for smaller values of β . We also note that, even for the input parameter set chosen by GM04, the degeneracy between the three parameters cannot be completely broken, though some parameter space can certainly be ruled out (see Fig. 7 of GM04). We consider that the redshift distribution only is not sufficient to measure all the three parameters by breaking the degeneracy completely.

4.1.2. The Type Ia Fraction

On the other hand, the difference of type Ia fraction is much larger than the redshift distribution in Fig. 4, indicating that this quantity is a better tool to constrain the parameters of interest. For the purpose of seeing this clearer, we made contour maps of the expected SN detection number and type Ia fraction, as a function of the two parameters of β and τ_{Ia} , which is shown in Fig 5. The contours of the detectable SN number and Ia fraction are mostly perpendicular to each other, demonstrating that they are good sets of observable quantities to break the degeneracy. Currently I_{AB} magnitude of ~ 24 is the spectroscopic limit for the largest ground-based telescopes, and it is difficult to measure the SN types beyond this magnitude. In this case the expected difference of Ia fraction is not very large ($f_{Ia} \sim 0.5$ – 0.6 for an interesting range of τ_{Ia}). On the other hand, future space borne spectroscopy or extremely large ground-based telescopes (30 m class) would allow the type classification at much deeper magnitudes. If SN types at $I_{AB} \sim 27$ are reliably determined, we expect that the Ia fraction differs by a factor of about 2 for the same range of τ_{Ia} .

In this analysis we have assumed that the effect of α on our predictions for shallow surveys is small. If the SN Ia fraction is reliably measured, the parameter β can be estimated without knowing α , since the core-collapse SN rate evolution to $z \sim 1$ simply reflects the SFR evolution. However, uncertainty of α affects the estimate of τ_{Ia} . We found that, when the α is varied from -2 to 1 , the estimate of τ_{Ia} by the same observational quantities is changed by 1–2 Gyr for the limiting magnitude of $I_{AB} = 24$ –27. This degeneracy can be broken by using deeper SN counts at magnitude of $\gtrsim 30$ (see below).

4.1.3. On the Form of the Ia Delay Function

The exact form of delay function is unknown, though there are some theoretical models (Yungelson & Livio 1998, 2000). Strolger et al. (2004) found that the Gaussian form of the delay function, $f_D(t) \propto \exp[-(t - \tau_{\text{Ia}})^2 / (2\sigma_{\text{Ia}}^2)]$ with $\sigma_{\text{Ia}} \sim (0.2\text{--}0.5)\tau_{\text{Ia}}$, fits better than the exponential form used so far in this paper. Therefore we repeated the above calculations but changing the delay function into the Gaussian form, to estimate the error on τ_{Ia} . It changes by about 1 Gyr, for the limiting magnitude of $I_{AB} \sim 27$. However, as shown by Strolger et al. (2004), the difference of the delay function appears most clearly in the redshift distribution of SNe, and hence the degeneracy with the delay function form may be removed by using the redshift distribution.

4.2. Determining α by SNe beyond $z \gtrsim 1$

It is obvious that we have to observe SNe well beyond $z \sim 1$ to determine α , and detection limit of $I_{AB} \sim 27$ is still not sufficient for this purpose (see Fig. 4). Future projects such as *SNAP* and *JWST* can detect supernovae down to ~ 30 magnitudes, and their data will be useful to constrain α , though it may be difficult to determine the SN types at such faintest magnitude.

We calculate the expected SN counts varying the α parameter and the results are shown in Fig. 6. In this calculation we choose a typical parameter set of $(\beta, \tau_{\text{Ia}}) = (2.0, 1.0)$. As expected, the change of SN counts by different α becomes larger for deeper observation. It should also be noted that the difference is much larger in the K band than I band, since the redshift distribution extends farther in redder bands ($z \sim 2$ for I and $z \gtrsim 5$ for K at ~ 30 mag). In a K -band observation with $m_{\text{lim}} = 30$, the fraction of SNe beyond $z = 2$ is 30–40 % in the case of $\alpha = 0$.

This is a calculation for fixed values of $(\beta, \tau_{\text{Ia}})$. By a supernova survey down to $I_{AB} \sim 27$ with secure type classification, the parameter β can be constrained well by using core-collapse SN statistics, while the degeneracy between τ_{Ia} and α remains. It is then possible to break the degeneracy by finding a set of $(\alpha, \tau_{\text{Ia}})$ satisfying both the type Ia statistics to $I_{AB} \sim 27$ and the all-type SN statistics to ~ 30 magnitudes.

4.3. Prospects for the Planned Projects

The current or past supernova surveys from ground, such as SCP (Pain et al. 2002) and HZT (Tonry et al. 2003), have reached a depth of about $I_{AB} \sim 24$. The recent GOODS project has achieved a depth of F850LP ~ 26 thanks to HST observations, but high signal-to-noise spectroscopic observation is still limited to $I_{AB} \sim$

24 by the largest ground telescopes. As shown above, the type Ia fraction needs to be determined with an accuracy distinguishing $\Delta f_{\text{Ia}} \sim 0.1$ to determine β and τ_{Ia} by surveys limited to $I_{AB} \sim 24$. Requiring statistical signal-to-noise of 5, a required number of SNe Ia becomes $N \gtrsim 1,000$, which is much larger than the number obtained so far. However, the ongoing ground-based projects such as SNLS (Pritchett et al. 2004) or ESSENCE (Garnavich et al. 2002; Matheson et al. 2004) will greatly increase the number of supernova to this magnitude. These surveys could give some useful constraint on β and τ_{Ia} , depending on the accuracy of f_{Ia} and systematic model uncertainties.

The situation would be improved if we can measure the Ia fraction down to $I_{AB} \sim 27$, as shown in Fig. 5. Spectroscopic type classification to this depth is impossible by current ground-based facilities, but classification by information obtained by imaging observations, such as light-curve shapes, colors, or host galaxy properties, may still be possible, as tried by Strolger et al. (2004) for the GOODS data. (See also Poznanski et al. 2002.) SNe II have very distinctive light-curves (type II-P or II-L), and are typically bluer than SNe Ia at early times. However, the reliability of this kind of approach is still rather uncertain so far; only about two thirds of the 42 GOODS SNe are secure type identifications (Dahlén et al. 2004). A major breakthrough in more distant future will be brought by spaceborne spectroscopy, such as *SNAP*, reliably measuring f_{Ia} to $I_{AB} \sim 27$. The *SNAP* will detect and get high signal-to-noise spectra of a few thousands of type Ia supernovae to $z \sim 1.7$ (Aldering et al. 2004), which corresponds to the peak magnitude of $I_{AB} \sim 27$ (see Fig. 4). Planned extremely large telescopes (~ 30 m class) will also be useful for type classification of the faintest SNe.

The spaceborne projects such as *SNAP* and *JWST* (Panagia 2003) will also allow us to break the degeneracy between α and τ_{Ia} by SN counts at magnitude of ~ 30 . As shown above, *SNAP* will have a sufficient number of supernovae but does not probe very high redshift beyond $z \sim 1$ because of the wavelength coverage. The *JWST* could probe higher redshift, but small field of view (14 square arcminutes) does not allow to detect many supernovae at one snapshot. According to our estimate with the parameter $\alpha = 0.0$, $\beta = 3.0$ and $\tau = 1.0$, *JWST* will find about 10 SNe down to $K \sim 30$ in one snapshot. To measure the supernova counts with an accuracy of ~ 10 –20% to obtain meaningful constraint on α , about 25–100 supernovae are required.

5. DISCUSSION AND CONCLUSION

In this paper we examined the dependence of observational quantities obtained by supernova surveys on the CSFH and the delay time of SNe Ia (τ_{Ia}) in detail, and tried to identify the best strategy to get meaningful constraints on them by planned future projects. Specifically, we chose three parameters of α , β , and τ_{Ia} , and investigated how well these parameters can be constrained in the future based only on supernova survey data, where α and β are the indices of the cosmic SFR evolution $[\propto (1+z)^{\{\alpha, \beta\}}]$ at $z \gtrsim 1$ or $\lesssim 1$, respectively.

Relatively shallow surveys ($I_{AB} \lesssim 27$) are relevant mainly with the parameters of β and τ_{Ia} . We have shown that it is essential to reliably measure the number frac-

tion of SNe Ia (f_{Ia}) in a survey sensitive to $I_{AB} \sim 27$, in order to measure these parameters. If a $\sim 20\%$ difference in f_{Ia} is clearly distinguished for supernovae at magnitude of $I_{AB} \sim 27$, a useful constraints on β will be obtained by core-collapse SN rate evolution. The constraint on τ_{Ia} is somewhat degenerate with the unknown α , but still it can be constrained within an accuracy of $\sim 1\text{--}2$ Gyr. Some regions of the parameter space of β and τ_{Ia} are seriously degenerated for the redshift distribution, and hence we consider that the Ia fraction is a better quantity for this purpose. However, the redshift distribution would be useful to constrain the functional form of the delay function (e.g., exponential or Gaussian).

Type classification down to $I_{AB} \sim 27$ may be potentially achieved by ongoing or near future ground-based surveys such as SNLS and ESSENCE, but type classification inevitably depends on less reliable methods based on imaging information. A more reliable study will become possible by spaceborne spectroscopy, as proposed for *SNAP*, or future extremely large telescopes on the ground. The parameter α can be constrained with breaking the degeneracy with τ_{Ia} , by ultra-deep SN surveys down to magnitudes of ~ 30 without SN type classification, which would become possible by future projects such as *SNAP* and *JWST*. The former has an advantage of larger statistics while the latter has an advantage of reaching higher redshifts by observing in longer wavelength. It should also be noted that, in the above discussion, we assumed that the local supernova rates, both for type Ia and core-collapse SNe, are well measured to give a good normalization that can be directly compared with high redshift observations. Therefore the effort to improve the accuracy of the local supernova rate is also highly encouraged (e.g. Aldering et al. 2002).

It should be noted that there is a number of simplifications in the calculation presented here, e.g., a constant IMF, no evolution in supernova light curves or spectra, a universal probability of supernova event occurrence per unit mass of star formation, and no evolution in the Ia delay time. These uncertainties might induce some bias in the CSFH or Ia delay time inferred from the SN rate history. On the other hand, it is possible to get information for these uncertain factors, if we use the CSFH determined by independent other methods (such as SFR estimates of high- z galaxies) as an external constraint.

Dust extinction has been incorporated in the same way for the SNe Ia and core-collapse SNe, with a fixed amount of extinction and no evolution with z . We adopted this simple treatment because it is quite difficult to construct a realistic model of dust extinction based on the present knowledge about galaxy evolution. However, it is likely that extinction has a wide distribution even within a galaxy (Hatano, Branch, & Deaton 1998), and mean extinction should also evolve with galaxy evolution (Totani & Kobayashi 1999). Because the time delay from star formation to supernova occurrence is different, core-collapse SNe could be much more heavily extincted by dust than SNe Ia (Mannucci et al. 2003). If the cosmic star formation is heavily hidden by dust at $z \gtrsim 2$ as inferred from submillimeter observations (e.g., Hughes et al. 1998), it might be appropriate to use different values of α for type Ia and core-collapse SNe. These issues should be kept in mind when the strategy discussed in this paper

TABLE 1
PEAK ABSOLUTE MAGNITUDES (VEGA), THEIR DISPERSION, AND
RELATIVE PROPORTIONS OF SUPERNOVA TYPES

Type	M_B	σ	f
Ia	-19.13	0.56 ^a	—
Ib/c	-17.71	1.39	0.23
IIP	-16.67	1.12	0.30
III	-17.70	0.90	0.30
IIn	-18.82	0.92	0.02

REFERENCES. — Richardson et al. (2002) for M_B and σ , and Dahlén & Fransson (1999) for f .

^aThe dispersion of SNe Ia is uncorrected about the light-curve-shape/luminosity relation.

is actually applied to future data. To see a typical effect of changing extinction, we repeated calculations with no extinction for SNe Ia and twice larger extinction for core-collapse SNe. The Ia delay time and β are then affected by ~ 1 Gyr and ~ 0.5 , respectively.

We presented our calculation of expected supernova counts as a function of variability flux in a given survey time interval, instead of the real flux. We also examined quantitatively the change of the detectable number of supernovae with time interval, and calculated the ratio of brightening and fading supernovae between the two epochs of the SN selection, which were not quantitatively reported in the earlier literature. These are useful for planning the observing intervals for the future observations. These quantities can easily be measured, and we examined a possibility of using these observational information to constrain the interesting parameters of CSFH and τ_{Ia} , but unfortunately they are not particularly useful. However, comparing these quantities to the theoretical predictions would be an important consistency check to examine, e.g., a possible contamination of other objects such as AGNs.

We would like to thank an anonymous referee for many constructive comments that greatly improved this paper. This work has been supported by the Grant-in-Aid for the 21st Century COE "Center for Diversity and Universality in Physics" and for TT (16740109) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.




FIG. 1.— Lightcurve and temperature templates for SNe of various types. The left panel is for the lightcurves used in our calculation and the right panel is temperature obtained by blackbody-fits to spectra. These are taken from SN 1999em (Hamuy et al. 2001), 1979C (Branch et al. 1981) and 1995G (Pastorello et al. 2002) for SN types of IIP, IIL and IIn. For Ib/Ic, these are constructed based on the data of 1994I, 1999ex and 2002ap (Richmond et al. 1996; Stritzinger et al. 2002; Foley et al. 2003). The data for SNe Ia is from Nugent et al. (2002).




FIG. 2.— Left : Expected number of SNe as a function of limiting magnitude of variability flux, in a field of 1 deg^2 . Various intervals of observation are assumed as indicated in the figure. The peak CSFH model and $\tau_{\text{Ia}} = 1.0 \text{ Gyr}$ are assumed (see text). Right : The ratio of expected SN number with an observing interval of 30 days to that of 360 days. Different line markings correspond to different models of CSFH (“peak” or “flat”) and the Ia delay time (0.3 or 3 Gyr), as indicated in the panel.




FIG. 3.— Left : The number ratio of brightening to fading SNe between the two epochs of supernova selection. Various intervals of observation are assumed as indicated in the figure. The peak CSFH model and $\tau_{Ia} = 1.0$ Gyr are assumed. Right : Different line markings correspond to different models of CSFH (“peak” or “flat”) and the Ia delay time (0.3 or 3 Gyr), as indicated in the panels. The time interval is 30 days.




FIG. 4.— Redshift distributions of SNe in a survey limited by $I_{AB} = 24$ and 27 , for two models (thick and thin curves) whose total expected SN number is almost the same but the CSFH parameters and type Ia delay time are largely different. The parameters are $(\beta, \tau_{Ia}[\text{Gyr}]) = (3.0, 4.0)$ for thick lines and $(2.3, 0.3)$ for thin lines. The solid and dashed lines show the distribution of total number including all SN types and that of SNe Ia, respectively.




FIG. 5.— Contour maps of the detectable SN number (thin curves) in a square-degree field and the fraction of SNe Ia (thick curves), for a survey with limiting magnitude of $I_{AB} = 24$ and 27. The x-axis is the delay time of SNe Ia (τ_{Ia}) and the y-axis is the low-redshift evolutionary index of the CSFH (β in Eq. 5). The search interval is assumed to be 30 days. If the interval is 360 days, the SN counts become about twice the values indicated in the figure, while the SN Ia fraction changes very little (at most by ~ 0.05).




FIG. 6.— Detectable SN number normalized by the case of $\alpha = 0$ as a function of limiting magnitude, for all SN types. Various α are assumed as indicated in the figure. The used parameters are $(\beta, \tau_{Ia}[\text{Gyr}]) = (2.0, 1.0)$. Left and right panels are for I and K' band, respectively.

REFERENCES

- Aldering, G., et al. 2004, preprint (astro-ph/0405232)
- Aldering, G., et al. 2002, *Proc. SPIE*, 4836, 61
- Blanc, G., et al. 2004, *A&A*, 423, 881
- Branch, D., Falk, S. W., Uomoto, A. K., Wills, B. J., McCall, M. L., & Rybski, P. 1981, *ApJ*, 244, 780
- Cappellaro, E., 2003, in *Supernova and Gamma-Ray burst*, ed. K. Weiler, (Heidelberg: Springer), 37
- Cappellaro, E., Evans, R., & Turatto, M. 1999, *A&A*, 351, 459
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Dahlén, T., & Fransson, C. 1999, *A&A*, 350, 349
- Dahlén, T., et al. 2004, *ApJ*, 613, 189
- Foley, R. J., et al. 2003, *PASP*, 115, 1220
- Furusawa, H., Shimasaku, K., Doi, M., & Okamura, S. 2000, *ApJ*, 534, 624
- Gal-Yam, A., Maoz, D., & Sharon, K. 2002, *MNRAS*, 332, 37
- Gal-Yam, A., Maoz, D., Guhathakurta, P., & Filippenko, A. V. 2003, *AJ*, 125, 1087
- Gal-Yam, A. & Maoz, D. 2004, *MNRAS*, 347, 942 (GM04)
- Garnavich, P. M. et al. 2002, *BAAS*, 34, 1233
- Gilliland, R. L., Nugent, P. E., & Phillips, M. M. 1999, *ApJ*, 521, 30
- Hamuy, M., et al. 2001, *ApJ*, 558, 615
- Hardin, D., et al. 2000, *A&A*, 362, 419
- Hatano, K., Branch, D., & Deaton, J., 1998 *ApJ*, 502, 177
- Hughes, D. H., et al. 1998, *Nature*, 394, 241
- Kobayashi, C., Tsujimoto, T., & Nomoto, K. 2000, *ApJ*, 539, 26
- Lanzetta, K. M., Yahata, N., Pascarelle, S., Chen, H., & Fernández-Soto, A. 2002, *ApJ*, 570, 492
- Lilly, S. J., Le Fevre, O., Hammer, F., & Crampton, D. 1996, *ApJ*, 460, L1+
- Madau, P., della Valle, M., & Panagia, N. 1998, *MNRAS*, 297, L17+
- Madau, P., Ferguson, H. C., Dickinson, M. E., Giavalisco, M., Steidel, C. C., & Fruchter, A. 1996, *MNRAS*, 283, 1388
- Madgwick, D. S., Hewett, P. C., Mortlock, D. J., & Wang, L. 2003, *ApJ*, 599, L33
- Mannucci, F., et al. 2003, *A&A*, 401, 519
- Maoz, D. & Gal-Yam, A. 2004, *MNRAS*, 347, 951
- Matheson, T., et al. 2004, preprint, (astro-ph/0411357)
- Nugent, P., Kim, A., & Perlmutter, S. 2002, *PASP*, 114, 803
- Pain, R., et al. 2002, *ApJ*, 577, 120
- Pain, R., et al. 1996, *ApJ*, 473, 356
- Panagia, N. 2003, preprint, (astro-ph/0309417)
- Pastorello, A., et al. 2002, *MNRAS*, 333, 27
- Perlmutter, S., et al. 1997, *ApJ*, 483, 565
- Phillips, M. M. 1993, *ApJ*, 413, 105
- Phillips, M. M., Lira, P., Suntzeff, N. B., Schommer, R. A., Hamuy, M., Maza, J. 1999, *AJ*, 118, 1766
- Poznanski, D., Gal-Yam, A., Maoz, D., Filippenko, A. V., Leonard, D. C., & Matheson, T. 2002, *PASP*, 114, 833
- Pritchett, C. J. for the SNLS collaboration, 2004, preprint, (astro-ph/0406242)
- Richardson, D., Branch, D., Casebeer, D., Millard, J., Thomas, R. C., & Baron, E. 2002, *AJ*, 123, 745
- Richmond, M. W., et al. 1996, *AJ*, 111, 327
- Schiminovich, D., et al., 2005, *ApJ*, 619, L47
- Stritzinger, M., et al. 2002, *AJ*, 124, 2100
- Strolger, L., et al. 2004, *ApJ*, 613, 200
- Sullivan, M., Treyer, M.A., Ellis, R.S., Bridges, T.J., Milliard, B.D., & Donas, J. 2000a *MNRAS*, 312, 442
- Sullivan, M., Ellis, R., Nugent, P., Smail, I., & Madau, P. 2000b, *MNRAS*, 319, 549
- Tonry, J. L., et al. 2003, *ApJ*, 594, 1
- Totani, T. & Kobayashi, C. 1999, *ApJ*, 526, L65
- Totani, T., Sato, K., & Yoshii, Y. 1996, *ApJ*, 460, 303
- Totani, T. and Takeuchi, T.T., *ApJ*, 570, 470
- Wilson, G., Cowie, L.L., Barger, A.J., Burke, D.J. 2002, *AJ*, 124, 1258
- Yungelson, L. & Livio, M. 1998, *ApJ*, 497, 168
- Yungelson, L. R. & Livio, M. 2000, *ApJ*, 528, 108